



International Scientific Organization
<http://iscientific.org/>
Current Science Perspectives
www.bosaljournals.com/csp/



Impact of effective microorganisms actuate (EMa) on development of Barley, Corn and Chard plants

Raghad S. Mouhamad^{1,*}, Ali S. Fadhel¹, Shaimaa A. Yousir¹, Ibrahim B. Razaq¹, Munawar Iqbal², Saba Ayub³ and Umer Younas²

¹Soil and Water Resources Center, Ministry of Sciences & Technology, Baghdad –Iraq

²Department of Chemistry, The University of Lahore, Lahore, Pakistan

³Govt. College University, Faisalabad, Pakistan

*Corresponding author's E-mail: raghad1974@yahoo.com

ARTICLE INFO

Article type:

Research Article

Article history:

Received July 2016

Accepted December 2016

January 2017 Issue

Keywords:

Microorganisms

Barley

Corn

Chard

ABSTRACT

An investigation trial was done at Agriculture Research Station, South Baghdad, Iraq, amid the two progressive periods of 2012 and 2013 to consider the impact of effective microorganisms (EM) at 5% on Barley, Corn and Chard, from the acquired results demonstrated that it is desirable over include EM at 5% as a dirt application isolated into one measurements connected at three times a season to improve synthetic properties of soil, biomass yield, and fiery debris, seed minerals substance of plants.

© 2017 International Scientific Organization: All rights reserved.

Capsule Summary: The impact of effective microorganisms (EM) on properties of soil as well as Barley, Corn and Chard biomass yield, and fiery debris, seed minerals substance was investigated.

Cite This Article As: Raghad S. Mouhamad, Ali S. Fadhel, Shaimaa A. Yousir, Ibrahim B. Razaq, Munawar Iqbal, Saba Ayub and Umer Younas. Impact of effective microorganisms actuate (EMa) on development of Barley, Corn and Chard plants. Current Science Perspectives 3(1) (2017) 60-66

INTRODUCTION

Powerful Microorganisms (EM) is a monetarily accessible fluid containing an assortment of lactic corrosive microbes, yeasts and phototrophic microscopic organisms (Iwaishi, 2000). These life forms make conditions which support common backing and empower them to outcompete hurtful pathogens (Shah et al., 2001), the various applications incorporated the accompanying: enhancing soil conditions for better plant development (Eissa, 2002), improving fertilizer creation and expanding the timeframe of realistic

usability of reaped products (Khaliq et al., 2006; Sahain et al., 2007).

Cultivating rehearses, accomplish manageability in horticultural generation, natural compost and bio-manure play a critical and key part. Since they have numerous alluring, soil properties and applies valuable impact on soil physical, substance and natural attributes (Boliłłowa and Gleń, 2008; Górski and Kleiber, 2010). Natural materials are utilized for expanding crop creation yet immaculate natural cultivating can never take care of the expanding demand for supplement supply, as adequate amounts of natural materials are not accessible (Frąszczak et al., 2011). Along these lines, one of the option of supplement supply is the joining of

Effective Microorganisms [EM] inoculums and natural/inorganic materials. It is proposed that Effective Microorganisms Activate (EMa) build the quantities of useful microorganisms in soil, in this way keeping up the characteristic biological community of the developed land and reducing the danger of ecological contamination with enhanced harvest efficiency and quality. Microorganisms improve the adequacy of culturing frameworks because of their part in disintegration of waste plant, harmonious and fermentative procedures. The utilization of EM in yield generation is either by direct application to soil, to natural matter spread on the dirt surface or to the plant. Be that as it may, it might use to treating the soil natural matter at an alternate area and this material connected to the dirt preceding planting. in which successful microorganisms (EMs) were utilized as a part of development of different types of rural products (cotton, maize, clear potatoes, rice, triticale, wheat) and also plant plants (rose, gerbera, apple, apricot) affirming their positive impact on most plants, while it neglected to exhibit a constructive outcome Higa (2003) and Wielgosz et al. (2010), There for a field investigation was done to decide the impact of EMa on development, produce segments of grain, maize, rough cultivars.

MATERIAL AND METHODS

EM1

EM1, obtained from Okinawa- Japan through AL-Anam for natural planting stock EM (Higa, 2003)

Preparation of secondary EM

The 10 litters of EM optional were set up by blending 0.5 litters of molasses with measure of water, then added water to achieve 9 litters, and supplemented 0.5 litters of powerful microorganisms; EMA, which delivered by Iraqi Ministry of Science and Technology, Soil and water asset focus. The past blend was kept in dull tanks to a vigorously age for one week

Table 1: Soil chemical analysis of the experimental site

Parameter	Soil texture and values
Texture	Clay loam
PH	7.5
Ec (dS/m)	2.03
O.M%	0.57
CEC (cmol/Kg)	23.6
CaCO ₃ gm/kg	140
Na gm/kg	15
K gm/kg	2.9
Ca gm/kg	54
Mg gm/kg	17
N gm/kg	0.19
P gm/kg	0.15

till pH is 3.5. The dirt was earth topsoil, pH 7.5. The aftereffects of pre-planting soil examination of the test field, have been exhibited in Table 2. The medicines including, grain, maize, coarse; which looked at for their Growth, compound properties furthermore use of EMa as take after: T1: Pre blossoming foliar utilization of EMa at a weakening rate of 5 (300 l/ha) for four times. T2: Control (no utilization of EMa).

The medications were duplicated three times in a RCBD with a plot size of 25 m² (5×5 m). All the agronomic operations aside from those under study, had kept uniform for every one of the medications for three seasons without culturing. Information on development and development segments were gathered utilizing standard strategies and dissected amid the 2-year study were search efficiency and nutritive worth as far as crisp and dry weight, and minerals of N,P, K, Ca, Mg,Na, Zn, Mn, Cu, and Fe. Scavenge profitability as far as dry matter was resolved a few slices every year when 10 to half of the populace is blooming. Proximate investigation was performed on ground tests of the dry weight and the strategies portrayed by Page (1982). Information were examined utilizing the SAS 2000 programming and means were isolated utilizing the LSD strategy (Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

Nitrogen (%)

Table 3, shows that every tried treatment altogether expanded the plant nitrogen substance of Plants as contrasted and the control treatment in three periods of study. In any case, the outcomes on the normal N plant of stem, and seed was altogether ($p < 0.05$) higher in the EMa treatment in grain -ash; grain -seed; corn ; Swiss Chard; grain -ash; grain -seed (0.5 ± 0.07 ; 1.92 ± 0.502 ; 0.92 ± 0.19 ; 3.22 ± 0.826 ; 0.85 ± 0.24 and 2.27 ± 0.19 individually), than in the control plot at three season.

Phosphorus (%)

Table 3, demonstrates that 5% EM gave a comparative and higher constructive outcome on leaf phosphorus content in the principal seasons contrasted and other season. Then again, 5% EM treatment gave the most astounding beneficial outcome on grain seed phosphorus in substance in the second season.

Potassium, Magnesium, Calcium, and Sodium (ppm)

Table 2 shows that all tried EM medicines prompted high beneficial outcome on leaf and seed potassium content than the control treatment in both seasons. By and large, EM treatment ended up being the most proficient medications in this worry. 5% EMa treatment in the principal season was most elevated critical in magnesium content in grain plants while EM supplemented medicines in the second season gave higher beneficial outcome on leaf calcium content in Swiss Chard plant. Table 2, demonstrates that EMa medications in the primary season gave most astounding calcium contrasted and second season in appear grain plant and EM medicines in

Table 2: Effect Effective Microorganisms Activate (EMa) added to plants on chemical content in plants at winter 2012, summer 2012 and winter 2013 seasons

	Winter 2012		Summer 2012		Winter 2013	
	barley -ash	barley -seed	corn	Swiss Chard	barley -ash	barley -seed
N						
cont.	0.42±0.04	1.98±0.46	0.89±0.011	2.75±0.284	0.80±0.22	2.04±0.21
EM (5%)	0.5±0.07	1.92±0.502	0.92±0.19	3.22±0.826	0.85±0.24	2.27±0.19
P						
cont.	0.035±0.006	0.37±0.028	0.26±0.15	0.086±0.047	0.10±0.02	0.31±0.05
EM (5%)	0.05±0.025	0.37±0.02	0.15±0.12	0.090±0.051	0.12±0.05	0.31±0.12
K						
cont.	12190±3139.5	8153.5±1829.28	18230±923.09	18276±8083.1	18276±8083.1	5179.6±462
EM (5%)	13539.8±475	7557.6±1199.9	18030±3492.2	26498.8±3	16030.1±2888.2	5298.7±459
Mg						
cont.	1847.5±153.6	2088.5±369.8	3638.4±2954	7548±1473.9	1639±388.6	1820±496.0
EM (5%)	1858.1±219.9	2549±226.7	3357.7±1533	7408.7±2165	1666±296.3	1712±178.7
Ca						
cont.	285.05±41.16	1347.5±400.9	4226±812.65	23101±25894	560.4±228.3	686.6±613.5
EM (5%)	345.4±131.7	1217.6±289.04	558.3±1327.5	8743.8±3637	653.5±600.3	367.2±191.7
Na						
cont.	2914±534.5	18115±671.7	575.4±162.21	68576±26604	24753.3±3131	1896.3±615
EM (5%)	2472±150.9	24558.33±9577	1117.7±769.6	58741±27000	22766.2±2233.2	1886.3±603
Cu						
cont.	3.05±0.216	4.38±0.212	7.71±1.94	11.76±5.0	8.03±2.24	6.84±0.49
EM (5%)	3.74±0.62	71.7±163.2	10.4±8.28	12.3±2.56	6.91±0.70	7.66±0.78
Mn						
cont.	1.59±0.318	1.59±0.31	9.73±15.9	189.0±82.2	4.7±5.31	2.92±4.74
EM (5%)	2.63±0.54	2.35±1.167	18.5±15.6	440.3±96.1	6.45±4.84	2.95±1.08
Zn						
cont.	11.72±0.53	32.15±5.81	20.69±2.52	14.39±7.44	47.08±8.41	60.2±9.91
EM (5%)	12.96±4.65	32.7±6.84	20.6±7.62	18.65±10.44	50.2±11.02	54.2±3.05
Fe						
cont.	4.95±1.61	4.33±0.96	564.8±91.3	695.1±210.2	397.9±236.6	274.6±240
EM (5%)	5.43±2.29	5.26±2.24	705.4±126.9	440.3±128.64	316±89.5	164.0±76.8

the late spring season gave comparative and higher beneficial outcome on leaf calcium content as contrasted and the control treatment. Likewise the aftereffects of grain plants examination showed a noteworthy on sodium content in first season contrasted with second season in appear treatment.

Cooper, Manges, Zinc and Iron (ppm)

In addition, Cu, Mn, Zn, and Fe substance of leaf, and grain was essentially ($p < 0.05$) higher in the EM fertilizer plot than in the control plot. Plants treated with the EM indicated critical contrasts when contrasted with the plants untreated at various season, while the statures of the plants became essentially higher from the control plants at summer season. However there was no noteworthy distinctive between the statures of plants (Corn and Swiss chard plant) treated with the EM. These outcomes are in concurrence with a few past

tests of gainful microorganisms adequacy on product plants. As indicated by our own particular writing review, crosswise over 36 autonomous studies and 26 plant species tried, helpful microorganisms (EM) soil application builds plant yield by 30 % (Schenck et al., 2009). though the microorganisms don't have any quantifiable effect on the plants (Javaid, 2010). We found a constructive outcome of advantageous microorganisms application on the development and advancement plant under zero-culturing framework, as did others (Megali et al., 2014).

Soil chemical properties

The long haul soil revisions brought about critical changes in soil synthetic properties (Figure 1, 2, 3, 4, 5). Soil Ec and pH were fundamentally ($p < 0.05$) lower in the three EMa plots than in the control plot at various season. Besides, soil pH




Fig. 1: Effect Effective Microorganisms Activate (EMa) added to plants on Ec content in soil at winter 2012, summer 2012 and winter 2013 seasons




Fig. 2: Effect Effective Microorganisms Activate (EMa) added to plants on pH content in soil at winter 2012, summer 2012 and winter 2013 seasons




Fig. 3: Effect Effective Microorganisms Activate (EMa) added to plants on Organic matter content in soil at winter 2012, summer 2012 and winter 2013 seasons.




Fig. 4: Effect Effective Microorganisms Activate (EMa) added to plants on Nitrogen content in soil at winter 2012, summer 2012 and winter 2013 seasons.




Fig. 5: Effect Effective Microorganisms Activate (EMa) added to plants on Phosphore content in soil at winter 2012, summer 2012 and winter 2013 seasons.




Fig. 6: Effect Effective Microorganisms Activate (EMa) added to plants on Biomass at winter 2012, summer 2012 and winter 2013 seasons.

was altogether ($p < 0.05$) lower in the EM expansion than in the control at winter season 2013. Soil natural matter, absolute N, and P substance was altogether ($p < 0.05$) higher in the two EMa plots for three seasons than in the control plot. Wellspring of supplement for the EMa originated from the sustenance supplied to the worms. EM compost indicated expanded in supplement when connected to soil. The impact of worm on the synthetic, physical and organic properties of compost have been appeared to impact supplement creation in soil. Living beings in EM assume part in yield creation. Research and field ponders in all landmasses have demonstrated that the immunization of EM society to the dirt/plant environment can enhance soil quality, soil wellbeing, and the development, yield, and nature of harvests (Benito et al., 2003). These outcomes are in concurrence with past tests of gainful microorganism's adequacy on expanded soil mineralization (Abdelhamid, 2004), soil group structure (Roca-Peréz et al., 2009).

Barley biomass

As appeared in Figure 6 the table underneath, the foliar use of EM brought about a huge increment in the biomass of grain in two seasons (winter 2012& 2013). In any case, the EM application at second season interim with 1:500 weakenings has given the most elevated biomass contrasted and first season. The most elevated crisp weight (< 10 t/h) of Barley was acquired with EM connected at 5% at winter 2013 intervals; the most noteworthy dry weight (< 3 t/h) of grain with 0% at second season interims.

The got results demonstrated that the Effective Microorganisms Activate (EMa) have noteworthy useful consequences for biomass of grain. Likewise the utilization of powerful microorganisms in harvest creation and demonstrate the capability of this innovation to decrease manure utilize and expand the yield and nature of products. This outcomes concurrence with general positive results stay of helpful microorganisms on product biomass and plant advancement have been accounted for (Goyal et al., 2005; Ibrahim, 2012).

CONCLUSIONS

This analysis was embraced to figure out whether the use of Effective Microorganisms Activate (EMA) would diminish the pH, EC. Numerous past studies have recommended that the utilization of EM has brought about a critical diminishment in solids taking care of, likewise one of the real component which enhances soil fruitfulness and expanding the accessibility of supplement components and thus influenced the plant development. Increment in the quantity of leaves with high chlorophyll content, leaf territory are normal events in plants that are furnished with appropriate sustenance and this can build the photosynthetic action of the plants. Increment in leaf territory and number of leaves ought to result to higher rates of photosynthesis henceforth expanded plant development. For plants, a high rate of net carbon digestion can bring about higher biomass gathering, favoring future development and proliferation.

CONCLUSIONS

This analysis was embraced to figure out whether the use of Effective Microorganisms Activate (EMA) would diminish the pH, EC. Numerous past studies have recommended that the utilization of EM has brought about a critical diminishment in solids taking care of, likewise one of the real component which enhances soil fruitfulness and expanding the accessibility of supplement components and thus influenced the plant development. Increment in the quantity of leaves with high chlorophyll content, leaf territory are normal events in plants that are furnished with appropriate sustenance and this can build the photosynthetic action of the plants. Increment in leaf territory and number of leaves ought to result to higher rates of photosynthesis henceforth expanded plant development. For plants, a high rate of net carbon digestion can bring about higher biomass gathering, favoring future development and proliferation.

REFERENCES

- Abdelhamid, M.T., Horiuchi, T., Oba, S., 2004. Composting of rice straw with oilseed rape cake and poultry manure and its effects on faba bean (*Vicia faba* L.) growth and soil properties. *Bioresource Technology* 93, 183–189.
- Benito, M., Masaguer, A., Moliner, A., 2003. Chemical and microbial parameters for the characterization of the stability and maturity of pruning waste compost. *Biology and Fertility of Soils* 37, 184–189.
- Boligłowa, E., Gleń, K., 2008. Assessment of effective microorganisms activity (EM) in winter wheat protection against fungal diseases. *Ecological Chemistry and Engineering A* 15(1-2), 23-27.
- Eissa, E.M., 2002. Use of Biostimulants in Activation of Soil Microflora for Yield and Fruit Quality Improvements of "Canino" Apricot. *Journal of Agricultural Research* 28, 354-364.
- Frąszczak, B., Kałużewicz, A., Krzesiński, W., Lisiecka, J., Spizewski, T., 2011. Effect of differential temperature and photoperiod on growth of *Ocimum basilicum*. *Żemdirbystė-Agriculture* 98(4), 375-382.
- Gomez, K.A., Gomez, A.A., 1984. Statistical procedure for agricultural research. 2nd (eds). John Wiley and Sons, Inc. New York.
- Górski, R., Kleiber, T., 2010. Effect of Effective Microorganisms (EM) on nutrient contents in substrate and development and yielding of Rose (*Rosa x hybrida*) and Gerbera (*Gerbera jamesonii*). *Ecological Chemistry and Engineering* 17(4), 505-513.
- Goyal, S., Dkull, S.K., Kapoor, K.K., 2005. Chemical and biological changes during composting of different organic wastes and assessment of compost maturity. *Bioresource Technology* 96, 1584–1891.
- Higa, T., 2003. EM Technology application in agriculture and environment protection. *Proceedings of the 38 International Microbiological Symposium Effective Microorganisms (EM) in Sustainable Agriculture and*

- Environmental Protection. SGGW, Rogów k/Łodzi, Poland. 17-18.
- Ibrahim, W. M. A., 2012. Behaviour of Taimour mango trees to inorganic and organic fertilization and application of EM. Ph. D. Thesis, Fac. of Agric. Minia Univ., Egypt.
- Javaid, A., 2010. Beneficial microorganisms for sustainable agriculture. In: Lichtfouse E (ed) Genetic engineering, biofertilisation, soil quality and organic farming. Sustainable agriculture reviews, 4. Springer, Netherlands, 347-369.
- Khaliq, A., Abbasi, M.K., Hussain, T., 2006. Effects of integrated use of organic and inorganic nutrient sources with effective microorganisms (EM) on seed cotton yield in Pakistan. *Bioresource Technology* 97, 967-972.
- Megali, L., Glauser, G., Rasmann, S., 2014. Fertilization with beneficial microorganisms decreases tomato defenses against insect pests. *Agronomy for Sustainable Development* 34, 649-656.
- Osman, S.O.A., Moustafa, F.M.A., Abd El-Galil, H.A., Ahmed, A.Y.M., 2011. Effect of yeast and Effective Microorganisms (Em1) application on the yield and fruit characteristics of Bartamuda date palm under Aswan conditions. *Assiut Journal of Agricultural Sciences* 42, 332-349.
- Roca-Peréz, L., Martínez, C., Marcilla, P., Buloda, R., 2009. Composting rice straw with sewage sludge and compost effects on the soil-plant system. *Chemosphere* 75, 781-787.
- Sahain, M.F.M., Abd El Motty, E.Z., El-Shiekh, M.H., Hagagg, L.F., 2007. Effect of some biostimulant on growth and fruiting of an apple trees in newly reclaimed areas. *Research journal of agriculture and biological sciences* 3(5), 422-429.
- Schenck_ Z.U., Schweinsberg-Mickan, M., Müller, T., 2009. Impact of effective microorganisms and other biofertilizers on soil microbial characteristics, organic-matter decomposition, and plant growth. *Journal of Plant Nutrition and Soil Science* 172(5), 704-712.
- Shah, H.S., Farrukh Saleem, M., Shahid, M., 2001. Effect of different fertilizers and Effective Microorganisms on growth, yield and quality of maize. *International Journal of Agriculture and Biology* 3(4), 378-379.
- Wielgosz, E., Dziamba, S.Z., Dziamba, J., 2010. Effect of application of EM spraying on the populations and activity of soil microorganisms occurring in the root zone of spring barley. *Polish Journal of Soil Science* 43, 1-12.
- Iwaishi, S., 2000. Effect of organic fertilizer and effective microorganisms on growth, yield and quality of paddy-rice varieties. *Journal of Crop Production* 3, 269-273.
- Page, A.L., Miller, R.H., Kenney, D.R., 1982. *Method of Soil Analysis*, 2nd (ed) Agron. 9, Publisher, Madiason, Wisconsin.

Visit us at: <http://bosajournals.com/csp/>

Submissions are accepted at: editorcsp@bosajournals.com